

``杏輝'

# **滴朗** 眼液 **DELONE** Eye Drop."**Simplar**"

本製劑乃配合具有強力消炎效能之Dexamethasone及具有廣效性抗菌作用之新黴素所成,故能有效治療眼部之疾患。

#### 成分:每1mL中含:

Dexamethasone sodium phosphate equivalent to Dexamethasone phosphate......1mg Neomycin sulfate......3.5mg(Potency)

賦形劑: Boric acid、Sodium borate、Sodium chloride、Menthol、Benzalkonium chloride、Water for injection

#### 滴應症:

, 健康敏感症、臉緣炎、非特異性表層結膜炎、深部角膜炎、 帶狀疱疹眼炎、水泡性角膜結膜炎、鞏膜角膜炎、結膜炎、 虹彩睫狀體炎、虹彩炎、復發性臉緣潰瘍、角膜潰瘍、及角膜損傷。

# 用法用量:

眼用-日間每小時一次,夜間每兩小時一次,每次滴一至二滴 於眼結膜囊内,如症狀減輕,可漸減為每四小時一滴,然後 再減為每日三至四次,每次一至二滴即可。

## 注意事項:

- 1.需有適當之抗菌治療,始可使用於細菌感染之病例。
- 2.若遇頑治之前眼部疾患或深部疾患時,須合併用腎上腺皮質荷爾蒙全身性治療。
- 3.一般遺傳性及退化性眼疾患,對本劑無效。
- 4.眼之結核性症及單純性疱疹乃本劑之禁忌症。
- 5.對於治療眼色素層炎,特別是慢性者,有時會引起眼壓高者,應予注意。
- 6.2歲以下兒童之使用須詢問醫師或藥師。

儲存:25℃以下保存。

本藥須由醫師處方使用

衛署藥製字第019915號

包裝:100毫升以下玻璃瓶裝、塑膠瓶裝。



® 杏輝藥品工業股份有限公司 SINPHAR PHARMACEUTICAL CO., LTD.

台灣 宜蘭 縣 冬 山 鄉 中 山 村 中 山 路8 4 號 84, CHUNG SHAN ROAD, CHUNG SHAN VILLAGE, TUNG - SHAN SHINE, I - LAN, TAIWAN 消費者服務專線: (0800)015151 website: http://www.sinphar.com

# **DELONE** Eye Drop. "Sinphar"

Delone combines the antiinflammatory activity of dexamethasone sodium phosphate with the antimicrobial activity of neomycin, thus permitting effective management of many disorders of the eye.

### Compositions: Each mL contains:

Dexamethasone sodium phosphate equivalent to Dexamethasone phosphate......1mg
Neomycin sulfate......3.5mg (potency)

Excipients:Boric acid 
Sodium borate 
Sodium chloride 
Menthol 
Benzalkonium chloride 
Water for injection

Indications: Lid allergy blepharitis, non-specific superficial keratitis, deep keratitis, herpes zoster ophthalmicus, phlytenular keratoconjunctivitis, keratoscleritis, conjunctivitis, iridocyclitis, iritis, recurrent marginal ulceration, keratohelcosis, corneal injury.

### Dosage and administration:

Instill one or two drops of ophthalmic solution into the conjunctival sac every hour during the day and every two hours during the night as initial therapy. When a favorable response is observed, reduce dosage to one drop every four hours. Later, further reduction in dosage to one drop three or four times daily many suffice to control symptoms.

#### Precautions:

- (1) If infection do not respond promptly, ophthalmic solution should be discontinued until the infection has been adequately controlled by other measures.
- (2) In stubborn cases of anterior segment eye disease and the deeper ocular structures are involved, systemic adrenocortical hormone therapy may be required.
- (3) Hereditary and degenerative eye disease in general do not show any response to treatment with this preparation.
- (4) It should not be used in the presence of infectious tuberculous lesions of the eye, herpes simplex.
- (5) By treatment of the acute anterior uvea, particulary chronic patient, may result in the intraocular pressure, must be carefully observed.

Packaging: Supplied in plastic bottle in less than 100mL.

